

# UC Davis

## UC Davis Previously Published Works

### Title

Draft Genome Sequence of the Caffeine-Degrading Methylophilum methylotrophicum populi  
Pinkel.

### Permalink

<https://escholarship.org/uc/item/1qk2q14t>

### Journal

Microbiology resource announcements, 8(47)

### ISSN

2576-098X

### Authors

Parales, Rebecca E  
Sharma, Gaurav  
Zhang, Xiangsheng  
et al.

### Publication Date

2019-11-01

### DOI

10.1128/mra.01300-19

Peer reviewed



# Draft Genome Sequence of the Caffeine-Degrading Methylotroph *Methylobacterium populi* Pinkel

Rebecca E. Parales,<sup>a</sup> Gaurav Sharma,<sup>a\*</sup> Xiangsheng Zhang,<sup>a,b</sup> Gabriel A. Subuyuj,<sup>a</sup> Jordan T. Langner,<sup>c</sup> Matthew E. Wright,<sup>a\*</sup> Jayna L. Ditty,<sup>c</sup>  Scott C. Dawson<sup>a</sup>

<sup>a</sup>Department of Microbiology and Molecular Genetics, College of Biological Sciences, University of California, Davis, Davis, California, USA

<sup>b</sup>Jiangsu Provincial Key Laboratory of Coastal Wetland Bioresources and Environmental Protection, Yancheng Teachers University, Yancheng, China

<sup>c</sup>Department of Biology, University of St. Thomas, St. Paul, Minnesota, USA

**ABSTRACT** A pink-pigmented facultative methylotroph, *Methylobacterium populi* Pinkel, was isolated from compost by selective enrichment with caffeine (3,5,7-trimethylxanthine) as the sole carbon, nitrogen, and energy source. We report here its high-quality draft genome sequence, assembled in 35 contigs totaling 5,630,907 bp. We identified 5,681 protein-coding sequences, including those putatively involved in caffeine degradation.

Caffeine (3,5,7-trimethylxanthine) is a natural purine alkaloid synthesized by many plant species. Most of the >35 caffeine-degrading bacterial isolates that have been reported to date are members of the genus *Pseudomonas* (1). We report here the isolation and draft genome sequence of a facultative methylotroph, *Methylobacterium populi* Pinkel, which can utilize caffeine as a sole carbon, nitrogen, and energy source, and predict that it uses the *N*-demethylation pathway to convert caffeine to xanthine.

Strain Pinkel was obtained from compost that contained coffee grounds, following enrichment in a variation of BG-11 minimal medium (2) lacking citrate and sodium nitrate, with 0.006 g/liter ferric citrate substituted for ferric ammonium citrate and containing 5 mM caffeine as the sole carbon and nitrogen source. A pink-pigmented pure culture was obtained after multiple passes on plates of the same medium. The strain was identified as *Methylobacterium populi* (3, 4) based on 16S rRNA gene amplification with universal 16S primers followed by sequence homology analysis. As expected for a *Methylobacterium* isolate, the strain was a rod-shaped facultative methylotroph capable of growing on methanol as sole carbon source (3, 5). It grew on 5 mM caffeine as the sole source of carbon, nitrogen, and energy with a doubling time of approximately 6.5 h.

Genomic DNA was purified using an ArchivePure DNA kit (5 Prime, Inc., Gaithersburg, MD). The genomic DNA (100 ng) was used to create a paired-end library using the Ovation Ultralow protocol (NuGen, Redwood, CA) by Illumina. Later, the library was sequenced using the Illumina MiSeq platform, producing 1,058,337 paired-end reads (length, 300 bp). Raw reads were quality trimmed with a minimum Phred quality score of 30 from both ends using Trimmomatic v0.35 (6) with default parameters, generating 973,280 (91.96% of total) sequencing reads of good quality. *De novo* assembly and scaffolding were performed using A5-miseq (version 20150522) (7), followed by the removal of duplicate contigs and contigs containing only repeat regions. Finally, contigs were cross-checked for any putative contamination using a Kraken-based taxonomic sequence classification system (8). The final assembly consists of 35 contigs (>500 bp) with a total size of 5,630,907 bp, an *N*<sub>50</sub> value of 589,043 bp, a G+C content of 69.2%, and ~40× coverage. The smallest and the largest contigs were 753 bp and 1,232,378 bp, respectively. Subsequently, the genome annotation was carried out with Rapid Annotations using Subsystems Technology (RAST) (9), which predicted a total of

**Citation** Parales RE, Sharma G, Zhang X, Subuyuj GA, Langner JT, Wright ME, Ditty JL, Dawson SC. 2019. Draft genome sequence of the caffeine-degrading methylotroph *Methylobacterium populi* Pinkel. Microbiol Resour Announc 8:e01300-19. <https://doi.org/10.1128/MRA.01300-19>.

**Editor** Steven R. Gill, University of Rochester School of Medicine and Dentistry

**Copyright** © 2019 Parales et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Rebecca E. Parales, [reparales@ucdavis.edu](mailto:reparales@ucdavis.edu), or Scott C. Dawson, [scdawson@ucdavis.edu](mailto:scdawson@ucdavis.edu).

\* Present address: Gaurav Sharma, Institute of Bioinformatics and Applied Biotechnology, Bengaluru, Karnataka, India; Matthew E. Wright, Benchling, San Francisco, California, USA.

**Received** 21 October 2019

**Accepted** 24 October 2019

**Published** 21 November 2019

5,885 coding sequences, including 5,681 protein coding sequences, 52 tRNA genes, 2 rRNAs (16S and 23S), and 150 repeat regions. Out of 5,681 proteins, 1,414 (~25%) of the genes were assigned to SEED subsystems, whereas 4,267 (~75%) were predicted to have an unknown (nonhypothetical) function. Subsystems signifying the survival of the isolate in aromatic compound-contaminated soil included membrane transport (90 genes), stress response (62 genes), metabolism of aromatic compounds (29 genes), and motility and chemotaxis (93 genes). As expected for a *Methylobacterium* species, the genome contained genes for carotenoid biosynthesis, flagellar motility and chemotaxis, and methanol utilization. In addition, genes for the *N*-demethylation pathway for caffeine degradation were also identified.

**Data availability.** This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [WEKV00000000](#). The version described in this paper is the first version, WEKV01000000. The data are under BioProject number [PRJNA514782](#) and BioSample number [SAMN10160078](#). The reads are available under SRA accession number [SRR8435744](#).

## ACKNOWLEDGMENT

*M. populi* Pinkel was isolated and characterized preliminarily by students at the University of California, Davis, during the 2014 Microbial Diversity Laboratory course (MIC 105L).

## REFERENCES

- Summers RM, Sridhar SK, Gopishetty S, Subramanian M. 2015. Genetic characterization of caffeine degradation by bacteria and its potential applications. *Microb Biotechnol* 8:369–378. <https://doi.org/10.1111/1751-7915.12262>.
- Rippka R, Waterbury JB, Stanier RY. 1981. Isolation and purification of Cyanobacteria: some general principles. In Starr MP, Stolp H, Trüper HG, Balows A, Schelgel HB (ed), *The prokaryotes*, vol 1. Springer-Verlag, Berlin, Germany.
- Green PN, Ardley JK. 2018. Review of the genus *Methylobacterium* and closely related organisms: a proposal that some *Methylobacterium* species be reclassified into a new genus, *Methylobacterium* gen. nov. *Int J Syst Evol Microbiol* 68:2727–2748. <https://doi.org/10.1099/ijsem.0.002856>.
- Van Aken B, Peres CM, Doty SL, Yoon JM, Schnoor JL. 2004. *Methylobacterium populi* sp. nov., a novel aerobic, pink-pigmented, facultatively methylotrophic, methane-utilizing bacterium isolated from poplar trees (*Populus deltoides* × *nigra* DN34). *Int J Syst Evol Microbiol* 4:1191–1196. <https://doi.org/10.1099/ijse.0.02796-0>.
- Green PN. 2005. Genus I: *Methylobacterium*, p 567–571. In Brenner DJ, Krieg NR, Staley JT (ed), *Bergey's manual of systematic bacteriology*, vol 2. Springer, New York, NY.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <https://doi.org/10.1093/bioinformatics/btu661>.
- Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. *Genome Biol* 15:R46. <https://doi.org/10.1186/gb-2014-15-3-r46>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, 3rd, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. *Sci Rep* 5:8365. <https://doi.org/10.1038/srep08365>.